

Nuclear Stress Test Instructions

	Patient Name	Medical Record #
Appointment Date	 Time	_ Lexiscan Adenosine Exercise

Note - Our office will call to verbally confirm your appointment the day before your test. Once confirmed, the technologist orders the radioisotope doses specially for you (which cannot be returned). If you are unable to make it to your scheduled appointment, please call 682-499-1777 **at least 24 hours ahead to avoid a \$150 fee** (which must be paid before scheduling your next appointment). If any insurance related complications with the pre-authorization process arise, our office will inform you ahead of time to reschedule your test.

MANDATORY: Please **BRING** a bagged **LUNCH TO THE TEST** for clear images. You will be instructed on when to eat by the technologist.

Requests:

- Please DO NOT bring children under the age of 10 to the appointment
- Please arrive **10 minutes early** for your scheduled appointment. If you are more than 15 minutes late, you will be rescheduled.

Preparing for the test:

- The entire test usually lasts about 4 hours
- Lexiscan patients please bring a caffeinated drink and a sandwich to have after the test
- DO NOT use any creams, lotions, ointments, oils, or powders over the chest or stomach area
- Wear comfortable clothing and walking shoes; women no dresses please
- If you use a **Nitro patch**, please **remove it 2 days prior** to your test
- Please bring a list of medications and/or actual bottles (including over the counter and herbal supplements). We need <u>complete</u> medication names & dosages
- If you use an inhaler, please bring it with you to your appointment
- Please inform if you are allergic to latex or rubber products, X-ray dye, or shellfish
- Please inform if you are pregnant, breast feeding, or think you may be pregnant

24 Hours Before

DO NOT take beta-blockers	ex: Sectral (Acebutolol), Tenormin (Atenolol), Zebeta (Bisoprolol), Lopressor & Toprol XL (Metoprolol), Corgard (Nadolol), Coreg (Carvedilol), Bystolic (Nebviolol), Inderal & InnoPran XL (Propranolo)	
NO Calcium Channel Blockers	ex: Norvasc (Amlodipine), Cardizem & Tiazac (Diltiazem), Felodipine, Isradipine, Nicardipine, Adalat CC & Procardia (Nifedipine), Sular (Nisoldipine), Calan & Verelan (Verapamil)	
NO Caffeine products after 8 pm the night before the test	ex: coffee, tea, chocolate, colas, energy drinks, some pain relievers (Anacin, Asprin, Excedrin), Theophylline, noncaffeinated, decaffeinated, and caffeine-free products(initial)	

4 Hours Before

DO NOT eat or drink anything after midnight for morning appointments.
 If your appointment is scheduled late in the afternoon, DO NOT eat or drink for at least 8 hours prior



Morning of the test

- **DO NOT take oral diabetes medications and blood pressure medication at home**. Please bring any diabetes medications and **all blood pressure medication** with you to take after the test.
- If you take insulin, take half the prescribed dose in the morning, and bring the other half with you
- Continue all other medicines as scheduled; you may take medicines with sips of water
- Lexiscan patients please bring a caffeinated drink and a sandwich to have after the test

After the test

- **Drink plenty of fluids for the next 48 hours** after the procedure to allow the remaining tracer to be flushed out completely
- You can continue your daily activities
- Dr. Lathia will discuss the test results (results are not given on the same day)

Nuclear Stress Test Waiver

I have read the above	information and understand my res	onsibilities and obliga	tions. I understand that if
I am unable to make i	t to my scheduled appointment, I an	responsible to inform	by calling 682-499-1777
at least 24-hours ahe	ead to avoid a \$150 fee. By signing	this form, I agree to co	mply with the above.
	X	X	
Date	Patient Signature	Witn	ess Signature